

June 2018 update

See below the latest updates for the current 19th issue, which are all incorporated in the PDF file of the Kourovka Notebook available at <https://kourovka-notebook.org>; (see also <http://math.nsc.ru/~alglog/19tkt.pdf> and <https://arxiv.org/pdf/1401.0300.pdf>).

***16.79.** Is it true that in any finitely generated AT -group over a sequence of cyclic groups of uniformly bounded orders all Sylow subgroups are locally finite? For the definition of an AT -group see (A. V. Rozhkov, *Math. Notes*, **40** (1986), 827–836)

A. V. Rozhkov

*No, it is not true (A. V. Rozhkov, in *Group theory and its applications*, Proc. XXII School-Conf. on Group Theory, Kuban' Univ., Krasnodar, 2018, 126–131 (Russian)).

***16.85.** Suppose that groups G, H act faithfully on a regular rooted tree by finite-state automorphisms. Can their free product $G * H$ act faithfully on a regular rooted tree by finite state automorphisms?

V. I. Sushchanskiĭ

*Yes, it can (M. Fedorova, A. Oliynyk, *Algebra Discrete Math.*, **23**, no. 2 (2017), 230–236).

***16.86.** Does the group of all finite-state automorphisms of a regular rooted tree possess an irreducible system of generators?

V. I. Sushchanskiĭ

*Yes, it does (Ya. Lavrenyuk, *Geometriae Dedicata*, **183**, no. 1 (2016), 59–67).

***19.40.** Does Thompson's group F (see 12.20) have the Howson property, that is, is the intersection of any two finitely generated subgroups of F finitely generated?

I. Kapovich

*No, it does not. Otherwise any subgroup of F would have this property. But the wreath product $\mathbb{Z} \wr \mathbb{Z}$, which does not have the Howson property (A. S. Kirkinskiĭ, *Algebra Logic*, **20**, no. 1 (1981), 24–36), can be embedded in F (S. Cleary, *Pacific J. Math.*, **228**, no. 1 (2006), 53–61). (D. Robertson, *Letter of 24 April 2018*.)

19.50. A finite graph is said to be integral if all eigenvalues of its adjacency matrix are integers.

*a) Let G be a finite group generated by a normal subset R consisting of involutions. Is it true that the Cayley graph $Cay(G, R)$ is integral?

b) Let A_n be the alternating group of degree n , let $S = \{(123), (124), \dots, (12n)\}$ and $R = S \cup S^{-1}$. Is it true that the Cayley graph $Cay(A_n, R)$ is integral?

D. V. Lytkina

*a) Yes, it is true (D. O. Revin, *Letter of 21 April 2018*; A. Abdollahi, *Letter of 3 May 2018*). Both proofs suggested are based on character theory; here is the second one. It suffices to show that the eigenvalues of $Cay(G, R)$ are rational, since the eigenvalues of a simple graph are algebraic integers. It is known that every eigenvalue of $Cay(G, R)$ has the form $\theta_\chi = \frac{1}{\chi(1)} \sum_{r \in R} \chi(r)$ for some complex irreducible character χ of G (implicit on pages 175–177 in P. Diaconis, M. Shahshahani, *Z. Wahrscheinlichkeitstheorie Verw. Gebiete*, **57** (1981), 159–179, see also Theorem 9 in M. Ram Murty, *J. Ramanujan Math. Soc.*, **18**, no. 1 (2003) 1–20). Since the value of any complex character on an involution is an integer, it follows that θ_χ is rational.

***19.75.** Let P be a finite 2-group of exponent 2^e such that the rank of every abelian subgroup is at most r . Is it true that $|P| \leq 2^{r(e+1)}$? This bound would be sharp (for a direct product of quaternion groups).

B. Sambale

*No, it is not true, as follows from the examples constructed in (A. Yu. Ol'shanskiĭ, *Math. Notes*, **23** (1978), 183–185). (A. Mann, *Letter of 23 April 2018*.)

***19.101.** The maximum length of a chain of nested centralizers of a group is called its c -dimension. Let G be a locally finite group of finite c -dimension k , and let S be the preimage in G of the socle of G/R , where R is the locally solvable radical of G . Is it true that the factor group G/S contains an abelian subgroup of index bounded by a function of k ?

A. V. Vasil'ev

*Yes, it is true (A. A. Buturlakin, [arXiv:1805.00910](https://arxiv.org/abs/1805.00910)).