

## October 2018 update

See below the latest updates for the current 19th issue, which are all incorporated in the PDF file of the Kourovka Notebook available at <https://kourovka-notebook.org>; (see also <http://math.nsc.ru/~alglog/19tkt.pdf> and <https://arxiv.org/pdf/1401.0300.pdf>).

For convenience of the readers all the updates since the first appearance of the 19th edition are also listed below; the newest ones added in October 2019 are marked **NEW**.

**NEW** \***12.79.** Suppose that  $a$  and  $b$  are two elements of a finite group  $G$  such that the function

$$\varphi(g) = 1^G(g) - 1_{\langle a \rangle}^G(g) - 1_{\langle b \rangle}^G(g) - 1_{\langle ab \rangle}^G(g) + 2$$

is a character of  $G$ . Is it true that  $G = \langle a, b \rangle$ ? The converse statement is true.

*S. P. Strunkov*

---

\*No, not always: a counterexample is given by  $G = A_4$ ,  $a = b = (123)$ . (S. V. Skresanov, *Letter of 20 July 2018*).

\***16.79.** Is it true that in any finitely generated  $AT$ -group over a sequence of cyclic groups of uniformly bounded orders all Sylow subgroups are locally finite? For the definition of an  $AT$ -group see (A. V. Rozhkov, *Math. Notes*, **40** (1986), 827–836)

*A. V. Rozhkov*

---

\*No, it is not true (A. V. Rozhkov, in *Group theory and its applications*, Proc. XXII School-Conf. on Group Theory, Kuban' Univ., Krasnodar, 2018, 126–131 (Russian)).

\***16.85.** Suppose that groups  $G, H$  act faithfully on a regular rooted tree by finite-state automorphisms. Can their free product  $G * H$  act faithfully on a regular rooted tree by finite state automorphisms?

*V. I. Sushchanskiĭ*

---

\*Yes, it can (M. Fedorova, A. Oliynyk, *Algebra Discrete Math.*, **23**, no. 2 (2017), 230–236).

\***16.86.** Does the group of all finite-state automorphisms of a regular rooted tree possess an irreducible system of generators?

*V. I. Sushchanskiĭ*

---

\*Yes, it does (Ya. Lavrenyuk, *Geometriae Dedicata*, **183**, no. 1 (2016), 59–67).

**NEW** \***17.108.** Is the group  $\langle a, b, t \mid a^t = ab, b^t = ba \rangle$  linear?

If not, this would be an easy example of a non-linear hyperbolic group. *M. Sapir*

---

\*Yes, this group is linear. Indeed, as noticed by M. Sapir, this group is the mapping torus of an irreducible atoroidal self-monomorphism of a free group; thus it is virtually special, and hence  $\mathbb{Z}$ -linear by Theorem B in (M. F. Hagen, D. T. Wise, *Duke Math. J.*, **165**, no. 9 (2016), 1753–1813). (M. F. Hagen, *Letter of 6 August 2018*).

\***19.40.** Does Thompson's group  $F$  (see 12.20) have the Howson property, that is, is the intersection of any two finitely generated subgroups of  $F$  finitely generated?

*I. Kapovich*

---

\*No, it does not. Otherwise any subgroup of  $F$  would have this property. But the wreath product  $\mathbb{Z} \wr \mathbb{Z}$ , which does not have the Howson property (A. S. Kirkinskiĭ, *Algebra Logic*, **20**, no. 1 (1981), 24–36), can be embedded in  $F$  (S. Cleary, *Pacific J. Math.*, **228**, no. 1 (2006), 53–61). (D. Robertson, *Letter of 24 April 2018*).

**\*19.50.** A finite graph is said to be integral if all eigenvalues of its adjacency matrix are integers.

a) Let  $G$  be a finite group generated by a normal subset  $R$  consisting of involutions. Is it true that the Cayley graph  $\text{Cay}(G, R)$  is integral?

b) Let  $A_n$  be the alternating group of degree  $n$ , let  $S = \{(123), (124), \dots, (12n)\}$  and  $R = S \cup S^{-1}$ . Is it true that the Cayley graph  $\text{Cay}(A_n, R)$  is integral?

*D. V. Lytkina*

---

\*a) Yes, it is true (D. O. Revin, *Letter of 21 April 2018*; see also the reference for part (b) below; A. Abdollahi, *Letter of 3 May 2018*). Both proofs suggested are based on character theory; here is the second one. It suffices to show that the eigenvalues of  $\text{Cay}(G, R)$  are rational, since the eigenvalues of a simple graph are algebraic integers. It is known that every eigenvalue of  $\text{Cay}(G, R)$  has the form  $\theta_\chi = \frac{1}{\chi(1)} \sum_{r \in R} \chi(r)$  for some complex irreducible character  $\chi$  of  $G$  (implicit on pages 175–177 in P. Diaconis, M. Shahshahani, *Z. Wahrscheinlichkeitstheorie Verw. Gebiete*, **57** (1981), 159–179, see also Theorem 9 in M. R. Murty, *J. Ramanujan Math. Soc.*, **18**, no. 1 (2003) 1–20). Since the value of any complex character on an involution is an integer, it follows that  $\theta_\chi$  is rational.

**NEW** \*b) Yes, it is true (W. Guo, D. V. Lytkina, V. D. Mazurov, D. O. Revin, *Preprint*, 2019, [arXiv:1808.01391](https://arxiv.org/abs/1808.01391)).

**\*19.75.** Let  $P$  be a finite 2-group of exponent  $2^e$  such that the rank of every abelian subgroup is at most  $r$ . Is it true that  $|P| \leq 2^{r(e+1)}$ ? This bound would be sharp (for a direct product of quaternion groups).

*B. Sambale*

---

\*No, it is not true, as follows from the examples constructed in (A. Yu. Ol'shanskiĭ, *Math. Notes*, **23** (1978), 183–185). (A. Mann, *Letter of 23 April 2018*.)

**\*19.101.** The maximum length of a chain of nested centralizers of a group is called its  $c$ -dimension. Let  $G$  be a locally finite group of finite  $c$ -dimension  $k$ , and let  $S$  be the preimage in  $G$  of the socle of  $G/R$ , where  $R$  is the locally solvable radical of  $G$ . Is it true that the factor group  $G/S$  contains an abelian subgroup of index bounded by a function of  $k$ ?

*A. V. Vasil'ev*

---

\*Yes, it is true (A. A. Buturlakin, [arXiv:1805.00910](https://arxiv.org/abs/1805.00910)).